

# Analysis of compounds in dichloromethane extractives for Sawara Falsecypress (*Chamaecyparis pisifera*) outer heartwood

NIU Jing<sup>1</sup>, LIU Zhi-Ming<sup>\*1</sup>, WANG Xiang-Ming<sup>2</sup>, XU You-Ming<sup>3</sup>, WANG Qing-Wen<sup>1</sup>

<sup>1</sup>Key Laboratory of Bio-based Material Science and Technology, Ministry of Education, P.R. China, Northeast Forestry University, Harbin 150040

<sup>2</sup>Forintek Canada Corp. Quebec, QC Canada G1P 4R4

<sup>3</sup>Huazhong Agricultural University, Wuhan 430070

**Abstract:** The chemical components of dichloromethane extractives for Sawara Falsecypress heartwood were analyzed with GC/MS except for basic chemical composition analysis for heartwood with Chinese standard method. 14 kinds of compounds were identified according to the computer compounds library data. The major compounds in dichloromethane extractives comprised of terpene and naphthalene derivatives. The experiments of antifungal effects of the dichloromethane extractive on *Aspergillus niger* were also carried out. The result showed that the dichloromethane extractive from Sawara Falsecypress has no or weak antifungal capability.

**Keywords:** *Chamaecyparis pisifera*; Sawara Falsecypress; Outer heartwood; Dichloromethane extractives; GC/MS; Antifungal

## Introduction

Sawara Falsecypress (*Chamaecyparis pisifera* (Sieb. Et Zucc.) Endl.) is derived from *Chamaecyparis formosensis* (Wang *et al.* 2003) and is an evergreen coniferous tree that grows mainly in central Japan. Sawara Falsecypress individuals grow in wet conditions on lower slopes and near mountain streams, and often occur in rocky depressions. This species not only forms mixed forests with other conifers such as *Chamaecyparis obtusa*, *Thujopsis dolabrata* and *Tsuga diversifolia*, but also mixes with deciduous broad-leaved trees, including *Fagus crenata* and *Quercus mongolica* (var. *grosseserrata*). The life history of Sawara Falsecypress in typical old-growth mixed forests generally involves the production of seeds through sexual reproduction of mature individuals (Zhao 1988). The species plays an important role in retaining soil and sand, coast defense and wild animal protection. Cultivation of introduced Sawara Falsecypress to China started in 1936. A few years later, it was then transplanted to other provinces involving Shandong, Liaoning, Guangxi, Yunnan and the middle and lower reaches of Changjiang River. In 1968, it was relocated from Lushan botanical garden in Jiangxi province to Jiuhua farmland in Hubei province. Generally, Sawara Falsecypress is an ideal tree species with the height of 50

meter and straight trunk, whose lumber is hard and durable. Furthermore, white sapwood and brown yellow heartwood are its main characteristics. On the other hand, it is a kind of better timber for making tools, building bridge, boat, vehicle crosstie and furniture owing to its superior property of resisting decay and slinky texture. It contains abundance of cellulose which makes it outstanding material for making paper.

Sawara Falsecypress, which is famous for appreciate value, has been cultivated widely in China. The cultivation of Sawara Falsecypress is not restricted by such environmental factors as climate and soil, and the livability could reach as high as 90% in the production with the skewer insert method (Wu 1983; Pan 1994; Wang 1990; Zhao 1988; Lin 1987; Yu 1993; Chen 1993). At present, there are broad researches on the Cupressaceae (*Chamaecyparis*) in foreign countries. However, the report on the use of *Chamaecyparis* species in natural durability is scarce. In Japan, the wood of *Chamaecyparis obtusa* is evaluated for the use value of constructing important buildings such as temples and shrines and is also considered to have hygienic properties for use as counter tops in sushi bars (Koyama *et al.*, 1997). Yoshixol (4,4-dimethyl-6-methylene-2-cyclohexen-1-one (C<sub>9</sub>H<sub>12</sub>O)) extracted from several species of *Chamaecyparis* has been shown to possess insecticidal activity. Termiticidal activity has been reported for the heartwood of *Chamaecyparis lawsoniana* (McDaniel 1989) and seed extracts of this species exhibited juveniling activity against the yellow mealworm beetle *Tenebrio molitor* (Jacobson *et al.* 1975). Antibacterial properties have been cited for *Chamaecyparis* (Johnson *et al.* 2001; Xiao *et al.* 2001; Yatagai and Nakatani 1994) and Debiaggi *et al.* (1988) reported that an ethanolic extract from the leaves of *Chamaecyparis lawsoniana* had antiviral activity against Herpes simplex virus type 2. Some reports showed that the extractives of methyl alcohol from Sawara Falsecypress leaves had antibacterial activity against *Staphylococcus aureus* and obtained several antibacterial fraction.

In addition, the reports on analysis of the extractives by gas chromatograph combined with mass spectrometry (GC-MS) are limited. The alkyl- and methoxy-phenolic content in wood

Foundation project: This work was supported by the Foundation of Postdoctoral Research of Heilongjiang and Innovation Fund of Northeast Forestry University (DLCX2004019)

Received date: 2007-03-25; Accepted date: 2007-06-07

©Northeast Forestry University and Springer-Verlag 2007

Electronic supplementary material is available in the online version of this article at <http://dxdoi.org/10.1007/s11676-007-0042-8>

Biography: Niu Jing (1983-), Female, Master Graduate, Northeast Forestry University

\*Corresponding author: (Email: zhimingliuwhy@126.com)

Responsible editor: Hu Yanbo

extractives were determined by the solid-phase microextraction-gas chromatography/mass spectrometry (SPME–GC/MS) (Verónica Pinon *et al.* 2007). The identification of the extractives from heartwood of Scots pine was performed using GC–MS, which gave good yields of the most important extractives: pinosylvin, pinosylvin monomethyl ether, resin acids and free fatty acids (Dag Ekeberg *et al.* 2006). The detailed composition of 70 underivatized wood extractive components presenting in quaking aspen were unequivocally identified by GC–MS method, forty-four compounds by retention time and mass spectral comparison with standards. An additional 26 chromatographic peaks were assigned to broad chemical class using retention time and mass spectra features (Fernandez 2001).

In china, the studies on Sawara Falsecypress are confined itself to the growth and regeneration, young cutting from germinating shoots, growth variation and regularity, wood basic density and resource utility. (Du 1999; Zhang *et al.* 2005; Lei *et al.* 2005; Shi and Xu *et al.* 2006 ).

On the basis of documents retrieved, the objective of this

study was to determine the basic chemical composition of outer heartwood of Sawara Falsecypress, and the antifungal activity of dichloromethane extractives against *Aspergillus niger*.

## Materials and methods

Sawara Falsecypress (SF) was harvested at the eastern hillside of 850 meter above sea level, where is a gradient of 20 degree and has a good forest nutrient. Sawara Falsecypress samples were cut into three logs above ground, marked A, B and C, respectively (Fig. 1). After air-drying 2 months at the lab in Northeast Forestry University, each part log was separately cut into about twenties disks, whose thickness was 5–7 cm. In the experiment, the lowest disk of log A heartwood was cut into 4 parts depending on 5 growing rings from external heartwood to pith, then sampled correspondent part as HB-A-4, HB-A-3, HB-A-2 and HB-A-1, respectively (Fig. 2).




Fig. 1 Scheme of SF stem

Fig. 2 Samples Scheme of Disk A

The selected sample, HB-A-4, was ground into wood powder passing through 2-mm sieves. Its moisture content (MC), cold-water and hot-water solubility, 1% sodium hydroxide solubility, alcohol-benzene solubility, nitric acid-alcohol cellulose and holocellulose, and acid-insoluble lignin content was respectively tested according to the Chinese standard method of fibrous raw material (GB/T2677.2-93; GB/T2677.4-93; GB/T2677.5-93; GB/T2677.6-94; GB/T2677.10-95; GB/T2677.8-1994) (1997). The reduplication occurred twice.

After its chemical composition analysis, the heartwood powder (HB-A-4), was extracted with hexane ( $n\text{-C}_6\text{H}_{14}$ ) for 24 h, and the extracted residue was continued to be extracted with dichloromethane ( $\text{CH}_2\text{Cl}_2$ ) for 24 h again (Liu *et al.* 2005). 0.04 g/ml dichloromethane extractive was tested by GC/MS 6890N-5973 insert (Agilent, USA). The instrument control parameters were listed as follows,

### Method comments

DB-17 column, 30 meters, 0.25 mm ID, 0.25 micron film, FID detector.

### Instrument control parameters

Injection volume: 1.0 microliters (syringe size: 10.0 microliters);  
splitting ratio: 1:20;  
Inlet temperature: 260°C; Gas: He; Constant flow: 1ml/min;  
Vacuum Compensation: On

### Oven program

Oven initial temperature: 40°C, Rate: 20°C /min  
Second temperature: 160°C, Retention time: 6 min, Rate: 5°C /min  
Final temperature: 240°C, Final time: 2 min

### MS Acquisition Parameters

Link inlet temperature: 280°C, EI: 70eV; Ion source temperature: 230°C

Acquisition Mode: Scan

Scan Parameters: Low Mass: 15 amu, High Mass: 260 amu

Then the antifungal effects of the extractive of dichloromethane against *Aspergillus niger* were tested. The method was

shown as follows: each plate was injected with 0.5 mL spore solution, which was prepared after the fungal was activated, and then adding 15mL PDA media to each plate sterilized. In order to make it well-distributed, the plate should be rotated sufficiently. Then 100  $\mu$ L of 0.06 g/ml dichloromethane extractive (acetone as solvent) was moved into the hole, which was made in the middle position of the plate. Compared to the one with extractive another two parallel experimental groups were assigned, one was injected with acetone and the other was blank. All the testing plates were incubated in 28°C for seven days, then observing the inhibition diameter of the fungi growth.

## Results and discussion

### Chemical Composition of Sawara Falsecypress


For wood powder of outer heartwood (HB-A-4), its moisture content, cold-water and hot-water solubility, 1% sodium hydroxide solubility, alcohol-benzene solubility, nitric acid-alcohol cellulose and holocellulose and acid-insoluble lignin content was respectively tested according to the Chinese standard method of fibrous raw material. The result in Table 1 was the average of twice measurements. These data were the basis of further study.

### Analysis of Dichloromethane Extractives Components for Sawara Falsecypress

Dichloromethane extractives of Sawara Falsecypress were tested by using GC/MS as mentioned above. Fig. 3 was the total ion chromatogram (TIC) of dichloromethane extractives for Sawara Falsecypress. 17 kinds of compounds were separated and 14 kinds of constituents were identified according to the computer compounds library data. The majority of the compounds were found. Their relative contents and match quality were listed in Table 1.

**Table 1. The chemical composition of Sawara Falsecypress**

Item	Heartwood (HB-A-4)	Item	Heartwood (HB-A-4)
MC%	8.60	Alcohol-benzene solubility%	4.61
Cold-water solubility%	6.60	Nitric acid-alcohol cellulose%	51.52
Hot-water solubility%	7.35	Holocellulose%	80.04
1% NaOH solubility%	15.47	Acid-insoluble lignin%	32.72



**Fig. 3 Total Ion Chromatogram of dichloromethane extractives for Sawara Falsecypress**

According to testing results, the major compound in dichloromethane extractives is 1,2,3,4,4a,5,6,8a-octahydro-7-methyl-4-methylene-1-(1-methylethyl)-, (1.alpha.,4a.alpha.,8a.alpha.)-naphthalene (Peak 14), its content is the highest (35.235%). But its match quality is lower, this compound need to be further identified by standard substance or other techniques such as elemental analysis and NMR analysis. The contents of Peak 11 and Peak 12 are a little lower compared with that of Peak 14, but higher than any other peaks, which their components are respective (+)-Epi-bicyclosquiphellandrene (13.291%) and alpha-Cadinol (14.143%). Generally, 14 kinds of compounds were

identified, major compounds comprised of terpenes and aromatic hydrocarbon.

According to the data in Table 1, the compounds of naphthalene derivatives were added up to 51.54%, about half of the total extracts. Meanwhile, a little vanillin existed in the extracts.

### Analysis of antifungal effects of Dichloromethane Extractives on *Aspergillus niger*

It is seen that there are no differences among the plates which the

*Aspergillus niger* grows as shown in Fig. 4, in other words, dichloromethane extractive has no or weak antifungal effects. According to the analysis of components in hexane extractive and dichloromethane one (Liu *et al.* 2006), both the highest contents are the same compound, namely 1,2,3,4,4a,5,6,8a- Octahydro-7-methyl-4-methylene-1-(1-methylethyl)-, (1.alpha.,4a.alpha., 8a.alpha.)-naphthalene. Dichloromethane extractive firstly extracted with hexane has fewer compounds than hexane one. In view of the variance, the antifungal tests in hexane extractive of Sawara Falsecypress can be carried out, then by virtue of the testing results, the antifungal components maybe exist in hexane extractive, and the antifungal components can be predicated, which is useful for further study. Furthermore, it is of great significance to choose a better isolation method in the future research for the extractive.




Fig. 4 Antifungal effect of dichloromethane extractive

Table 2. The assignments of peaks for the TIC of dichloromethane extractives in Fig.3.

Peak No.	Retention Time/min	Compounds	Relative Content%	Match Quality
1	10.841	Naphthalene,1,2,3,4,4a,5,6,8a-octahydro-7-methyl-4-methylene-1-(1-methylethyl)-,(1.alpha.,4a.alpha.,8a.alpha.)	0.931	99
2	11.221	unknown	1.497	–
3	11.547	Naphthalene,1,2,4a,5,6,8a-hexahydro-4,7-dimethyl-1-(1-methylethyl)-	3.568	97
4	11.616	Naphthalene,1,2,3,5,6,8a-hexahydro-4,7-dimethyl-1-(1-methylethyl)-,(1S-cis)-	4.958	95
5	12.071	Naphthalene,1,2,3,4-tetrahydro-1,6-dimethyl-4-(1-methylethyl)-,(1S-cis)-	1.209	96
6	12.435	Vanillin	0.947	96
7	12.602	Benzene,1-methyl-3-[(1-methylethylidene)cyclopropyl]-	1.162	72
8	13.338	Naphthalene,1,2,3,4,4a,7-hexahydro-1,6-dimethyl-4-(1-methylethyl)-	1.081	78
9	13.589	unknown	2.716	–
10	13.733	Naphthalene,1,2,4a,5,6,8a-hexahydro-4,7-dimethyl-1-(1-methylethyl)-	0.853	62
11	13.938	(+)-Epi-bicyclosquiphellandrene	13.291	90
12	14.029	alpha.-Cadinol	14.143	91
13	14.181	Copaene	6.575	97
14	14.371	Naphthalene,1,2,3,4,4a,5,6,8a-octahydro-7-methyl-4-methylene-1-(1-methylethyl)-,(1.alpha.,4a.alpha.,8a.alpha.)-	35.235	60
15	17.801	Cedren-13-ol,8-	3.756	89
16	20.002	2-Naphthalenemethanol,decahydro-.alpha.,.alpha.,4a-trimethyl-8-methylene-, [2R-(2.alpha.,4a.alpha.,8a.beta.)]-	3.614	62
17	24.366	unknown	4.464	–

Conclusion

The moisture content, cold-water and hot-water solubility, 1% sodium hydroxide solubility, alcohol-benzene solubility, nitric acid-alcohol cellulose and holocellulose, and acid-insoluble lignin content was respectively tested for wood powder of Sawara Falsecypress heartwood. These data showed the basic chemical composition of Sawara Falsecypress heartwood. It is concluded that the 1,2,3,4,4a,5,6,8a-octahydro-7-methyl-4-methyene-1-(1-methylethyl)-, (1.alpha.,4a.alpha.,8a.alpha.)-naphthalene is the major component of dichloromethane extractives for Sawara Falsecypress heartwood. Its relative content is the highest (35.235%), but its match quality is lower. Thus this compound

needs to be further identified by standard substance or other techniques such as elemental analysis and NMR analysis. Another 2 kinds of compounds, whose contents are higher, are (+)-Epi-bicyclosquiphellandrene (13.291%) and alpha-Cadinol (14.143%), respectively. The major compound in dichloromethane extractives comprised of terpene and naphthalene derivatives. These compounds have no antifungal capability.

References

Cai Cheng, Zhang Fengzhi, *et al.* 2005. Test Study on Propagation by Cutting with *Chamaecyparis pisifera*. *Hubei Forestry Science and Technology*, 5: 23–24.(in Chinese)  
Chen Zilin, Sun Lingxian. 1993. Growth Study of *Chamaecyparis pisifera* cv.

- plumosa in Pan'an Town. *Journal of Zhejiang Forestry College*, **10**(4): 424–426. (in Chinese)
- Chinese Light Industry Standard of Paper Making. 1997. GB /T 2677. 2, GB /T 2677. 4, GB /T 2677. 5, GB /T 2677. 6, GB /T 2677. 8, GB /T 2677. 10
- Dag Ekeberg, Per-Otto Flåte, Morten Eikenes, *et al.* 2006. Qualitative and quantitative determination of extractives in heartwood of Scots pine (*Pinus sylvestris* L.) by gas chromatography. *Journal of Chromatography*, **1109**: 267–272.
- Debiaggi, M., Pagani, L., Cereda, P.M., *et al.* 1988. Antiviral activity of *Chamaecyparis lawsoniana* extract: study with *Herpes simplex* virus type 2. *Microbiologica*, **11**: 55–61.
- Du Youxin. 1999. A study on the better wide-ringed species of *Chamaecyparis pisifera*. *Jiangxi Forestry Science and Technology*, **3**: 9–10. (in Chinese)
- Fernandez. M. P., Watson. P. A., and BreuilGas C. 2001. Chromatography-mass spectrometry method for the simultaneous determination of wood extractive compounds in quaking aspen. *Journal of Chromatography*, **922**: 225–233.
- Hayakawa Takehiro, Tomaru Nobuhiro, Yamamoto Shin-ichi. 2004. Stem distribution and clonal structure of *Chamaecyparis pisifera* growing in an old-growth beech-conifer forest. *Ecological Research*, **19**: 411–420.
- Jacobson, M., Redfern, R.E., Mills, G.D. 1975. Naturally occurring insect growth regulators part 2 screening of insect and plant extracts as insect juvenile hormone mimics. *Lloydia*, **38**(6): 455–472.
- Johnson, A.P., Aucken, H.M., Cavendish, S., *et al.* 2001. Dominance of EMRSA-15 and -16 among MRSA causing nosocomial bacteraemia in the UK: analysis of isolates from the European Antimicrobial Resistance Surveillance System (EARSS). *Journal of Antimicrobial Chemotherapy*, **48**: 143–144.
- Koyama, S., Yamaguchi, Y., Tanaka, S., *et al.* 1997. A new substance (Yoshixol) with an interesting antibiotic mechanism from wood oil of Japanese traditional tree (Kiso-Hinoki), *Chamaecyparis obtusa*. *General Pharmacology*, **28**(5): 797–804.
- Lei Xiaohua, Zhang Fengzhi, Chen Shuangtian, *et al.* 2005. Growth variation of *Chamaecyparis pisifera* under different site condition. *Hubei Forestry Science and Technology*, **6**: 17–20. (in Chinese)
- Lin Xie. 1987. Fundamental study on growth regularity of *Chamaecyparis pisifera* and *Chamaecyparis obtusa*. Forestry Technology communication, (12): 10–13. (in Chinese)
- Liu Zhiming, Niu Jing, Wang Xiangming, *et al.* 2006. Analysis of compounds in hexane extractives of Sawara Falsecypress heartwood. *Biomass Chemical Engineering*, **40**: 25–28. (in Chinese)
- Liu Zhiming, Xu Youming, Niu Jing, *et al.* 2005. Analysis of chemical composition for Sawara Falsecypress (*Chamaecyparis pisifera*). In Proceedings of International Symposium on Research & Effective Utilization of Bio-based Materials. Nanning, Guangxi, China, Dec. 2<sup>nd</sup>–4<sup>th</sup>: 628–631.
- Maruyama E., Hosoi Y., Ishii K. 2003. Somatic embryo culture for propagation, artificial seed production, and conservation of sawara cypress (*Chamaecyparis pisifera* Sieb. et Zucc.). *Journal of Forest Research*, **2**: 1–8.
- McDaniel, C.A., 1989. Major termiticidal components of heartwood of Port-Orford cedar, *Chamaecyparis lawsoniana* (A. Murr.). *Parl. Material und Organismen*, **24**(1): 1–16.
- Pan Zhigang, You Yingtian. 1994. *Transplanting and cultivating of the major exotic species in China*. Beijing: Beijing Science and Technology Press.
- Umezawa T. 2000. *Chemistry of extractives*. In: Shiraishi N, Hon DN-S (eds), Wood and cellulosic chemistry, 2<sup>nd</sup> ed. New York: Marcel Dekker, pp 213–241.
- Verónica Pino, Juan H. Ayala, Venerando González *et al.* 2007. Determination of the alkyl- and methoxy-phenolic content in wood extractives by micellar solid-phase microextraction and gas chromatography–mass spectrometry. *Talanta*, In Press, Corrected Proof, Available online 24 .
- Wang W P, Hwang C Y, Lin T P, *et al.* 2003. Historical biogeography and phylogenetic relationships of the genus *Chamaecyparis* (Cupressaceae) inferred from chloroplast DNA polymorphism. *Plant Systematics and Evolution*, **241**: 13–28.
- Wang Yueying, Jin Chuan. 1990. Fundamental study on transplanted experiment for 24 kinds of cypress. *Zhejiang Forestry Science and Technology*, **10**(2): 24–28, 38. (in Chinese)
- Wu Zonglun. 1983. *Introduction to exotic species*. Beijing: Science Press.
- Xu Youming, Shi Yuhu, Xu Zixue, *et al.* 2006. Growth regularity of *Chamaecyparis pisifera* and variations for latewood percentage and wood basic density. *J. of Northeast Forestry University*, **34**(1): 48–51. (in Chinese)
- Xiao, D., Kuroyanagi, M., Itani, T., *et al.* 2001. Studies on constituents from *Chamaecyparis pisifera* and antibacterial activity of diterpenes. *Chemical and Pharmaceutical Bulletin*, **49**(11): 1479–1481.
- Yatagai, M., Nakatani, N., 1994. Antimite, antitfly, antioxidative and antibacterial activities of pisiferic acid and its congeners. *Mokuzai Gakkaishi*, **40**: 1355–1362.
- Yu Ciyang. 1993. Fundamental study on transplanted sea island experiment for Virginia pencil cedar. *Zhejiang Forestry Science and Technology*, **13**(6): 16–21. (in Chinese)
- Zhai Hongwu. 2005. Cultivation technique of *Chamaecyparis pisifera*. *Agriculture of Henan*, **9**: 23. (in Chinese)
- Zhao Yinan. 1988. Introduction to cupressus species. *Anhui Forestry Science and Technology*, (3): 24–26. (in Chinese)